



Comparison of Susceptibility Status of laboratory and field populations of *Aedes aegypti* against Temephos in Rawalpindi

Dr. Ali Arslan, Dr. Uzair Mukhtar, . Shumaila Mushtaq, Arqam Bakhtiyar Zakki,
Muhammad Hammad and Adil Bhatti

Medical Entomology & Disease Vector Control (MEDVC), Health Services Academy, Islamabad

*Corresponding author: aliarslan761@gmail.com

Abstract

Background: In Pakistan, application of the larvicide temephos to the aquatic breeding sites of *Aedes aegypti* is a key part of the dengue control strategy. Throughout Pakistan no data is available on the susceptibility status of *Aedes aegypti* to temephos. The aim of the study was to examine the susceptibility status of *Aedes aegypti* against temephos and compare and estimate the resistance ratios of both laboratory and field populations on the basis of median lethal concentration LC_{50} and LC_{95} . **Methodology:** Larval bioassays were carried out according to WHO method on the field population of *Aedes aegypti* from three different localities (Rawal Town, Pothohar Town and Cantonment) of Rawalpindi. The laboratory reared susceptible strain of *Aedes aegypti* provided by the insectarium of Medical Entomology and Disease Vector Control (MEDVC) department was used as reference for comparison. **Results:** The LC_{50} values of all field populations of *Aedes aegypti* were recorded as 0.024 mg/L for Rawal Town, 0.019 mg/L for Pothohar Town and 0.025 mg/L for Cantonment showing resistance ratios RR_{50} of 6.32, 5 and 6.58 respectively. Similarly LC_{95} values were recorded as 0.26 mg/L for Rawal Town, 0.23 mg/L for Pothohar Town and 0.25 mg/L for Cantonment showing resistance ratios RR_{95} of 9.29, 8.21 and 8.93 respectively. **Conclusion:** Our results indicated that all the field populations of *Aedes aegypti* from Rawal Town, Pothohar Town and Cantonment were resistant to temephos, compromising the field effectiveness of this organophosphate. This comparative data of resistance ratios RR_{50} and RR_{95} on the temephos resistance of *Aedes aegypti* could help public health services to design more effective vector control measures.

Keywords: *Aedes aegypti*, insecticide susceptibility status and temephos.

Introduction

Dengue is an arboviral disease. It is caused by any one of the four serotypes (DEN-1, DEN-2, DEN-3 and DEN-4). These four distinct serotypes are closely related to each other (1). It is transmitted by the bites of mosquitoes of the genus *Aedes* and mostly by the species *Aedes aegypti* and *Aedes albopictus* etc. In the last 50 years, there is a 30 fold increase in the incidence of this disease. In the past decade, it has expanded from urban to rural areas (2).

According to World Health Organization (WHO), there are currently 50-100 million cases every year. Case fatality rate of dengue is about 2.5% (1).

The first confirmed outbreak of dengue fever (DF) was reported from Karachi in 1994. The epidemics

occurred continuously since 2005 in Karachi (3). There were about 3305 confirmed cases of DF in Karachi in 2010 (4). The disease remained endemic in the southern parts of Sindh and Baluchistan since 1994 however from 2004-5, dengue has spread across the country (5). Pakistan is now in endemic situation for DF in Khyber Pakhtun Khwa (KPK), Punjab and Sindh (3). The Swat district of KPK province in Pakistan has experienced an outbreak of DF in 2013. Since 7 August, suspected cases reached to a level of 6376 which also included 23 deaths having the death rate of about 0.36%. The outbreaks of DF continued to increase after the floods in 2010 (6). This epidemic situation has resulted in the 16580 confirmed cases of DF with 257 deaths in Lahore. In rest of the country there were approximately 5000 cases and 60 deaths

(3). In 2011, there were about 252935 suspected and 17057 laboratory confirmed cases with 219 deaths in Pakistan (7). Rawalpindi is facing an epidemic of dengue having the highest number of cases after Lahore in Punjab. It has the border areas with Islamabad and the Swat district of Khyber Pakhtun Khwa (KPK). It is facing more threat to dengue due to huge breeding places and increased number of cases in Islamabad and the recent outbreak of dengue in the Swat district of KPK.

Aedes aegypti is highly anthropophilic with markedly endophilic and endophagic behaviors; these characteristics are directly related to its high efficiency as a disease vector (8, 9). In the absence of a vaccine or effective therapeutic medications, the best strategy to control and prevent the disease is to control the vector populations (9). Control strategies are based on community participation which includes the removal of water from the containers in domestic and peri-domestic areas and environmental sanitation. These actions may also be supplemented with other vector control interventions such as the chemical control. Many dengue vector control interventions target the immature stages of the mosquito, which breed in artificial containers in close proximity to human dwellings. The most widely used method for controlling immature *Aedes aegypti* is the periodic treatment of actual and potential breeding sites with chemical larvicides. The organophosphate (OP) insecticide temephos is commonly used to control immature dengue vectors due to its cost-effectiveness and community acceptance (8, 10, and 11). The application of chemical insecticides is a major tool in minimizing the transmission of DF in most Latin American countries (12, 13). Resistance to this temephos has also been recorded in *Aedes aegypti* in Asia including Cambodia (14), Thailand (15, 16), and Malaysia (17) and South America (18, 19).

Temephos is the most widely used larvicide in our country and in Rawalpindi. However, its use is strictly subjected to available information of resistance to the local *Aedes aegypti* populations. This changing trend of resistance in target vector species needs to be properly monitored and assessed against the most widely used larvicide temephos. Current study was carried out to examine the susceptibility status of *Aedes aegypti* mosquito larvae against temephos in field populations of Rawalpindi and compare and estimate the resistance ratios of both laboratory and field populations on the basis of median lethal concentration (LC_{50}) and LC_{95} .

Materials and Methods

Mosquito Sampling:

The study was conducted in three different localities of Rawalpindi i.e. Rawal Town, Pothohar Town and Cantonment, based on the previous dengue endemic areas and areas having the insecticidal selection pressure. The larvae were collected from potential breeding sites of *Aedes* mosquito. For each sampling site, larvae from random breeding places were collected and stored in plastic boxes. These field collected larvae were then transferred to Medical Entomology and Disease Vector Control (MEDVC) insectarium of Health Services Academy, Islamabad for rearing to the adult stage.

Mosquito Rearing:

Field collected mosquitoes were reared to generation F1 for carrying out the larval bioassays. Mosquito populations were maintained at insectaries conditions ($27^{\circ}\text{C} \pm 2^{\circ}\text{C}$; relative humidity $80\% \pm 10\%$) and females were fed on albino mice to complete their gonotrophic cycle (20). While the susceptible reference strain of *Aedes aegypti* were provided by the Medical Entomology and Disease Vector Control (MEDVC) insectarium of Health Services Academy.

Preparation of Stock Solution:

1% stock solution was prepared by mixing 200 mg of active ingredient of temephos (0.4 ml of 50% E.C Temeguard^R) in 20 ml distilled water. Subsequent serial dilutions were made as per WHO guidelines. Wide range of concentrations (0.0001-0.5 PPM) was made as per WHO guidelines. A total of 100 larvae were exposed to each concentration of temephos (21).

Larval Bioassays:

Larval bioassay was conducted separately on both laboratory reared and field collected larvae. Batches of 25 late third or early fourth instar larvae were transferred by means of droppers to small disposable test cups, each containing 99 ml of water and 1 ml of temephos concentration. Small, unhealthy or damaged larvae were removed and replaced (21). Each bioassay included a control group which received 1 mL of ethanol. Tests were run at $27^{\circ}\text{C} \pm 2^{\circ}\text{C}$, and mortality was assessed after 24 h of insecticide exposure (21, 22).

Data from all the replicates was pooled and subjected to computer software programs (Minitab v 16.0) for analysis. LC₅₀ and LC₉₅ values were calculated from a log dosage–probit mortality regression line. Bioassays were repeated three times, using new solutions or suspensions and different batches of larvae each time. Resistance ratios (RR₅₀ and RR₉₅) of each population compared to the laboratory strain of *Aedes aegypti* (used as reference) were calculated as follows: RR₅₀ = LC₅₀ assay/LC₅₀ reference; RR₉₅ = LC₉₅ assay/LC₉₅ reference (22). If the control mortality is between 5% and 20%, the mortalities of treated groups are corrected according to Abbott’s formula (23).

A mosquito population was considered susceptible when RR₅₀ was less than 2, less resistant when RR₅₀ was between 2 and 5, moderately resistant when RR₅₀ was between 5 to 10 and highly resistant when RR₅₀ was over 10 (24).

Results

Percentage mortality of laboratory and field populations of *Aedes aegypti* larvae at various concentrations (mg/L) after 24 hours exposure is shown in table 1. The LC₅₀ value of laboratory reference strain was recorded as 0.0038mg/L while the LC₅₀ values from field populations of Rawal Town, Pothohar Town and Cantonment were recorded as 0.024 mg/L, 0.019 mg/L and 0.025 mg/L respectively. The LC₅₀ values from the field populations of Rawal Town, Pothohar Town and Cantonment were far more than the LC₅₀ value of laboratory susceptible reference strain. The LC₅₀ values of both laboratory

and field populations were pooled for analysis to estimate the resistance ratios (RR₅₀). The resistance ratio RR₅₀ was recorded as 6.32 for Rawal Town, 5 for Pothohar Town and 6.58 for Cantonment showing moderate resistance according to the criteria of Mazzari & Georghiou 1995 (24).

The LC₉₅ value of laboratory strain was recorded as 0.028 mg/L while the LC₉₅ values from field populations of Rawal Town, Pothohar Town and Cantonment were recorded as 0.26 mg/L, 0.23 mg/L and 0.25 mg/L respectively. The LC₉₅ values from the field populations of Rawal Town, Pothohar Town and Cantonment were far more than the LC₉₅ value of laboratory reference strain. The LC₉₅ values of both laboratory and field populations were pooled for analysis to estimate the resistance ratios (RR₉₅). The resistance ratio RR₉₅ was recorded as 9.29 for Rawal Town, 8.21 for Pothohar Town and 8.93 for Cantonment showing moderate resistance according to the criteria of Mazzari & Georghiou 1995 (24).

The estimated LC₅₀ and LC₉₅ of temephos for the laboratory susceptible reference strain and the three field populations of *Aedes aegypti* are presented in Table 2 and graph 1 and 2 where data of temephos lethal concentrations and resistance ratios are estimated. By analyzing the lethal concentrations, all the three field populations differed from the laboratory susceptible reference strain and also among each other. The results suggested that all the three field populations were moderately resistant to temephos.

Table 1: Percentage mortality of laboratory and field populations of *Aedes aegypti* larvae at various concentrations (mg/L) after 24 hours exposure:

Sample	Control	No. of dead larvae at various conc. (mg/L) post exposure 24 hours							
		0.0001	0.0005	0.001	0.005	0.01	0.05	0.1	0.5
Laboratory Population	13/600	0	0	59	114	172	225	225	225
Mortality (%)	2.17	0	0	26.22	50.67	76.44	100	100	100
Rawal Town Population	8/600	0	0	11	32	62	99	225	225
Mortality (%)	1.33	0	0	4.89	14.22	27.55	44	100	100
Pothohar Town Population	5/600	0	0	14	41	80	116	225	225
Mortality (%)	0.83	0	0	6.22	18.22	35.55	51.55	100	100
Cantonment Population	7/600	0	0	59	114	172	225	225	225
Mortality (%)	1.17	0	0	4	12.44	25.78	44	100	100

Table 2: Estimated lethal concentrations of temephos (Confidence interval) and resistance ratios based on the laboratory susceptible reference strain

Sample	Regression line	Pearson ² goodness of fit	LC ₅₀ (95% CI)	RR ₅₀	LC ₉₅ (95% CI)	RR ₉₅
<i>Aedes aegypti</i>						
Laboratory Population	0.82±0.037	53.02	0.0038 (0.0034-0.0042)	1	0.028 (0.023-0.035)	1
Rawal Town Population	0.69±0.029	136.34	0.024 (0.021-0.027)	6.32	0.26 (0.21-0.34)	9.29
Pothohar Town Population	0.66±0.028	111.7	0.019 (0.017-0.022)	5	0.23 (0.18-0.30)	8.21
Cantonment Population	0.72±0.031	132.33	0.025 (0.022-0.028)	6.58	0.25 (0.20-0.32)	8.93

LC₅₀: Lethal concentration which kills 50% of population expressed in mg/L

LC₉₅: Lethal concentration which kills 95% of population expressed in mg/L

CI: Confidence Interval

RR: Resistance ratio

RR₅₀: Resistance ratio 50%: LC₅₀ field strain/LC₅₀ lab strain

RR₉₅: Resistance ratio 95%: LC₅₀ field strain/LC₉₅ lab strain

P > 0.05 suggests a well-fitting model

P < 0.05 suggests an invalid model population

Figure 1: LC₅₀, LC₉₅ values of temephos against laboratory and field populations of *Aedes aegypti* larvae





Figure 2: RR₅₀, RR₉₅ values of temephos against laboratory and field populations of *Aedes aegypti* larvae

Discussion

This study was carried out to examine the susceptibility status of field populations of *Aedes aegypti* larvae against temephos in three different localities of Rawalpindi city i.e. Rawal Town, Pothohar Town and Cantonement. The larvicide temephos is the most commonly used chemical in Rawalpindi for the suppression of *Aedes aegypti* larvae in dengue control programs. These programs have often been implemented without information on the resistance selection risk posed by temephos in the field. The lack of data on *Aedes aegypti* susceptibility to temephos is a limiting factor for the success of control programs. For this reason, evaluations are often performed using the susceptibility of laboratory colonies as a reference, which do not necessarily reflect the natural variations that can be found among field populations, which has been demonstrated in other studies (25). The widespread use of temephos in controlling the *Aedes aegypti* larval population has led to resistance in many Latin American countries including Cuba (26), El Salvador (27), Argentina (28), Bolivia (29), Venezuela (30), Peru (31) and Colombia (32). It is believed that the extent of temephos resistance is underestimated due to under-reporting and lack of surveillance (11).

Our estimations of lethal concentrations LC₅₀ of laboratory reference strain of *Aedes aegypti* were

almost equal with the lethal concentrations of susceptible Rockefeller strain observed by Marcombe et al. (33) (LC₅₀ of 0.0037 mg/L); Rodriguez et al. (34) (LC₅₀ of 0.00364 mg/L). Our estimations of lethal concentrations LC₅₀ of laboratory reference strain of *Aedes aegypti* were lower than the lethal concentrations of susceptible Rockefeller strain observed by Uthai et al. (35) (LC₅₀ of 0.0060 ppm); Ponlawat et al. (36) (LC₅₀ of 0.0054 mg/L); and higher than the LC₅₀ observed by Andrighetti et al. (37) (LC₅₀ of 0.0025 ppm); Prophiro et al. (38) (LC₅₀ of 0.00167 mg/L); Bisset et al. (39) (LC₅₀ of 0.0012 mg/L); Gomez et al. (40) (LC₅₀ of 0.003058 mg/L). Our estimations of lethal concentrations LC₉₅ were slightly higher than the lethal concentrations observed by Araujo et al. (41) (LC₉₅ of 0.011 mg/L); Lima et al. (42) (LC₉₅ of 0.017mg/L). Our estimations of lethal concentrations LC₅₀ of field populations were slightly higher than the LC₅₀ of field populations observed by Singh, R. K in Delhi, India. The resistance ratios in all the field populations of Rawalpindi showed moderate to high resistance according to the Criteria of Mazzari & Georghiou 1995 (24). This high resistance pattern observed in the *Aedes aegypti* field populations of Rawalpindi may be associated with the prolonged use of temephos in each locality with different intensities or application frequencies thus subjecting them to different selection pressures with consequent increase in resistance mechanisms in each population.

Conclusion

The results of the present study confirm the detection of resistance in field population of *Aedes aegypti* larvae to temephos in Rawalpindi. The findings of temephos resistant larvae of *Aedes aegypti* field populations in Rawalpindi confirm the need for monitoring the susceptibility of these populations as one of the strategies for ensuring the efficacy of control programs for this vector in the area. Detection of resistance could help public health personnel to formulate appropriate alternative steps to ensure the effectiveness of control effort that may accompany with the emerging problems of insecticide resistance. Furthermore, cross resistance or resistance as a result of agricultural uses of insecticides may evolve and adversely impact the options to switch to an alternative method or insecticides for disease control. Our observations could help to guide insecticide-based strategies, although broader monitoring of insecticide resistance in *Aedes* mosquitoes is needed.

References

- World Health Organization. Dengue Fact sheet, September 2013 From URL <http://www.who.int/mediacentre/factsheets/fs117/en/index.html>
- World Health Organization, Dengue, from URL <http://www.emro.who.int/health-topics/dengue/index.html>
- World Health Organization, Pakistan, Programme areas, Dengue fever From URL <http://www.emro.who.int/pak/programmes/dengue-fever.html>
- Shamim M. Frequency, pattern and management of acute abdomen in dengue fever in Karachi, Pakistan. *Asian Journal of Surgery*. 2010; 33(3):107-13.
- Directorate of Malaria Control, National Guidelines for Dengue Vector (s) Control in Pakistan, Ministry of NHSRC, Islamabad, 2013.
- Saleem T, Saleem S. Knowledge, attitudes and practices regarding dengue fever among adults of high and low socioeconomic groups. 2010.
- World Health Organization, Dengue fever in Pakistan, September 2013 From URL <http://www.emro.who.int/surveillance-forecasting-response/outbreaks/dengue-fever-in-pakistan.html>
- World Health Organization, Dengue guidelines for diagnosis, treatment, prevention and control, new edition 2009.
- Scott TW, Morrison AC. Vector dynamics and transmission of dengue virus: implications for dengue surveillance and prevention strategies. *Dengue Virus*: Springer; 2010; 115-28.
- Chavasse D, Yap H. Chemical methods for the control of vectors and pests of public health importance. 1997.
- Ranson H, Burhani J, Lumjuan N, Black IV WC. Insecticide resistance in dengue vectors. *TropIKA.net*. 2010; 1(1).
- Georghiou G, Wirth M, Tran H, Saume F, Knudsen A. Potential for organophosphate resistance in *Aedes aegypti* (Diptera: Culicidae) in the Caribbean area and neighboring countries. *Journal of medical entomology*. 1987; 24(3):290-4.
- Gratz N. Emergency control of *Aedes aegypti* as a disease vector in urban areas. *Journal of the American Mosquito Control Association*. 1991; 7(3):353.
- Polson KA, Curtis C, Chang M, Olson JG, Chantha N, Rawlins S. Susceptibility of two Cambodian population of *Aedes aegypti* mosquito larvae to temephos during 2001. *Dengue Bull*. 2001;25:79-83
- Chareonviriyahpap T, Aum-Aung B, Ratanatham S. Current insecticide resistance patterns in mosquito vectors in Thailand. *The Southeast Asian journal of tropical medicine and public health*. 1999; 30(1):184-94.
- Paeporn P, Komalamisra N, Deesin V, Rongsriyam Y, Eshita Y, Thongrungrat S. Temephos resistance in two forms of *Aedes aegypti* and its significance for the resistance mechanism. *Southeast Asian journal of tropical medicine and public health*. 2003; 34(4):786-92.
- Chen C, Nazni W, Lee H, Sofian-Azirun M. Susceptibility of *Aedes aegypti* and *Aedes albopictus* to temephos in four study sites in Kuala Lumpur City Center and Selangor State, Malaysia. *Tropical biomedicine*. 2005; 22(2):207-16.
- Llinás GA, Seccacini E, Gardenal C, Licastro S. Current resistance status to temephos in *Aedes aegypti* from different regions of Argentina. *Memórias do Instituto Oswaldo Cruz*. 2010; 105(1):113-6.
- Macoris MdLG, Andrighetti MTM, Takaku L, Glasser CM, Garbeloto VC, Bracco JE. Resistance of *Aedes aegypti* from the state of São Paulo, Brazil, to organophosphates insecticides. *Memórias do Instituto Oswaldo Cruz*. 2003; 98(5):703-8.

20. Practical Manual and Guidelines for Dengue Vector Surveillance, Medical Research Institute and Coordination Unit, 2011.
21. World Health Organization, Guidelines for Laboratory and Field Testing of Mosquito Larvicides, 2005. (WHO/CDS/WHOPES/GCDPP/2005.13)
22. Kamgang B, Marcombe S, Chandre F, Ntchoutpouen E, Nwane P, Etang J, et al. Insecticide susceptibility of *Aedes aegypti* and *Aedes albopictus* in Central Africa. *International Pest Control*. 2011; 53(3):153.
23. Abbott WS. A method of computing the effectiveness of an insecticide. *Journal of Economic Entomology*, 1925; 18:265–67.
24. Mazarri MB, Georghiou GP. Characterization of resistance to organophosphate, carbamate and pyrethroid insecticides in field populations of *Aedes aegypti* from Venezuela. *Journal of Mosquito Control Association*. 1995; 11: 315-322.
25. Robertson J, Preisler H, Ng S, Hickie LA, Gelernter W. Natural variation: a complicating factor in bioassays with chemical and microbial pesticides. *Journal of Economic Entomology*. 1995; 88(1):1-10.
26. Bisset JA, Rodríguez MM, Fernández TD, Pérez O. Status of insecticide resistance and resistance mechanisms in larvae Playa municipality, collected during the intensive phase against *Aedes aegypti* in Havana, Institute of Tropical Medicine, 2001-2002. 2004; 56(1):61-6.
27. Bisset Lazcano JA, Rodríguez MM, San Martín JL, Romero JE, Montoya R. Assessing the insecticide resistance of an *Aedes aegypti* strain in El Salvador. *Revista Pan Americana de Salud Pública*. 2009; 26(3):229-34.
28. Llinás GA, Seccacini E, Gardenal C, Licastro S. Current resistance status to temephos in *Aedes aegypti* from different regions of Argentina. *Memórias do Instituto Oswaldo Cruz*. 2010; 105(1):113-6.
29. Biber PA, Dueñas JR, Almeida FL, Gardenal CN, Almirón WR. Laboratory evaluation of susceptibility of natural subpopulations of *Aedes aegypti* larvae to temephos. *Journal of the American Mosquito Control Association*. 2006; 22(3):408-11.
30. Rodríguez MM, Bisset J, de Fernandez DM, Lauzán L, Soca A. Detection of insecticide resistance in *Aedes aegypti* (Diptera: Culicidae) from Cuba and Venezuela. *Journal of medical entomology*. 2001; 38(5):623-8.
31. Rodríguez MM, Bisset JA, Fernandez D. Levels of insecticide resistance and resistance mechanisms in *Aedes aegypti* from some Latin American countries. *Journal of the American Mosquito Control Association*. 2007; 23(4):420-9.
32. Grisales N, Poupardin R, Gomez S, Fonseca-Gonzalez I, Ranson H, Lenhart A. Temephos Resistance in *Aedes aegypti* in Colombia Compromises Dengue Vector Control. *PLoS neglected tropical diseases*. 2013; 7(9):e2438.
33. Marcombe S, Darriet F, Agnew P, Etienne M, Yp-Tcha M-M, Yébakima A, et al. Field efficacy of new larvicide products for control of multi-resistant *Aedes aegypti* populations in Martinique (French West Indies). *The American journal of tropical medicine and hygiene*. 2011; 84(1):118-26.
34. Rodríguez, M. M., Bisset, J., Ruiz, M., & Soca, A. Cross-resistance to pyrethroid and organophosphorus insecticides induced by selection with temephos in *Aedes aegypti* (Diptera: Culicidae) from Cuba. *Journal of medical entomology*. 2002; 39(6):882-888.
35. Uthai L, Rattanapreechachai P, Chohanadisai L. Bioassay and effective concentration of Temephos against *Aedes aegypti* larvae and the adverse effect upon indigenous predators: *Toxorhynchites splendens* and *Micronecta* sp. *Asia Journal of Public Health*. 2011; 2(2).
36. Ponlawat A, Scott JG, Harrington LC. Insecticide susceptibility of *Aedes aegypti* and *Aedes albopictus* across Thailand. *Journal of Medical Entomology*. 2005;42(5):821-5
37. Andrighetti MTM, Cerone F, Rigueti M, Galvani KC, Macoris MdLdG. Effect of pyriproxyfen in *Aedes aegypti* populations with different levels of susceptibility to the organophosphate temephos. 2008.
38. Prophiro JS, Silva OS, Luna JED, Piccoli CF, Kanis LA, Silva MAND. *Aedes aegypti* and *Aedes albopictus* (Diptera: Culicidae): coexistence and susceptibility to temephos, in municipalities with occurrence of dengue and differentiated characteristics of urbanization. *Revista da Sociedade Brasileira de Medicina Tropical*. 2011; 44(3):300-5.
39. Bisset J, Marín R, Rodríguez M, Severson D, Ricardo Y, French L, et al. Insecticide resistance in two *Aedes aegypti* (Diptera: Culicidae) strains from Costa Rica. *Journal of medical entomology*. 2013; 50(2):352-61.

40. Gómez A, Seccacini E, Zerba E, Licastro S. Comparison of the insecticide susceptibilities of laboratory strains of *Aedes aegypti* and *Aedes albopictus*. Memórias do Instituto Oswaldo Cruz. 2011; 106(8):993-6.
41. Araújo AP, Diniz DFA, Helvecio E, De Barros RA, De Oliveira CMF, Ayres CFJ, et al. The susceptibility of *Aedes aegypti* populations displaying temephos resistance to *Bacillus thuringiensis israelensis*: a basis for management. Parasit Vectors. 2013; 6:297.
42. Lima EP, Paiva M, De Araújo AP, Da Silva E, Da Silva UM, De Oliveira LN, et al. Insecticide resistance in *Aedes aegypti* populations from Ceará, Brazil. Parasit Vectors. 2011; 4(5):2-12.
43. Singh, R. K., Mittal, P. K., Kumar, G., & Dhiman, R. C. Insecticide susceptibility status of *Aedes aegypti* and *Anopheles stephensi* larvae against temephos in Delhi, India. *International Journal of Mosquito Research*. 2014.