



2348-8069

DOI: <http://dx.doi.org/10.22192/ijarbs.2021.08.04.008>

Monthly and Daily Activity Pattern of Tsetse Fly *Glossina pallidipes* in Maze National Park, Gamo Zone, Southwest Ethiopia

Dr Abenezer Mara Zuma*

Melo Gada woreda livestock and fishery office, Gofa Zone, SNNPR, Ethiopia

E-mail: abenezerzuma2020@gmail.com

Abstract

The general pattern of fluctuations in tsetse populations is roughly related to the ambient and rainfall distribution. *Glossina pallidipes* inhabiting Maze National park was studied from November 2018 to May 2019 to investigate their seasonal and daily activity pattern. Acetone plus cow urine baited biconical traps were used to evaluate the seasonal activity pattern by taking sample every month for five consecutive days for five months and daily activity pattern was investigated by taking sample every hour for three consecutive days each in the months of November to March. The density of *G. pallidipes* was found to be high during wet months and decreased during dry months. *Glossina pallidipes* showed a bi-modal daily activity pattern with morning peak followed by mid-day depression and then late afternoon peak. Its activity rhythm showed a dual relation with temperature, positively correlated in the morning when the temperature is less than 36°C ($r = 0.699$ and $p = 0.122$) and negatively correlated when temperature is greater than 36°C ($r = -0.649$ and $p = 0.023$). The relationship between fly activity and relative humidity was also negatively correlated in the morning ($r = -0.649$, $P=0.1626$) and positively correlated in the afternoon ($r = 0.989$, $P = 0.002$). In general *G. Pallidipes* were more abundant during the rainy season and active around noon and in the evening. To improve agricultural development and livestock production, strategic tsetse control methods particularly the use of traps should follow the activity pattern of the vectors.

Keywords: Daily Activity, Maze National Park, *Glossina Pallidipes*, Seasonal Activity

Introduction

Tsetse flies (*Glossina* species) can be ranked among the world's most destructive pests and are the vectors of the causative agents for sleeping sickness in humans and African Animal Trypanosomosis (AAT) or Nagana in livestock [1]. The tsetse flies are adapted to the wide range of habitats, from central African humid rain forests to the vast semiarid, open savannahs of eastern Africa. The total area infested by tsetse flies is between 8.5 and 10 million km², which is more than 40% of the total land area of the infested countries [2]. Out of the nine regions of Ethiopia five

(Amhara, Beneshangul- Gumus, Gambella, Oromiya and SNNPR) are infested by more than one species of tsetse flies [3].

The general pattern of fluctuations in tsetse populations is roughly related to rainfall distribution throughout the year with a basic feature of an increase when rains start and a decrease during the dry season. Their pattern of distribution is a result of mainly climatic and biological factors [4]. For most species, the optimum temperature is about 25°C, and higher or lower temperatures during particular seasons or development stages, limit their existence [5]. Long

cold winters, often with frost, make conditions that are not suitable for tsetse. In cold weather, the fly cannot move about to feed. In cold soil, the pupa cannot complete its development before its stored food supplies are finished. The occurrence of tsetse flies requires the presence of their host. Therefore, insufficient food may cause some areas to be without tsetse, even though the areas may be suitable in other respects [6].

Tsetse fly exacts a high public health burden and has a devastating impact on livestock and agriculture. One of such disease that has plagued sub-Saharan Africa is caused by protozoan African trypanosomes (the trypanosome species) and transmitted by tsetse flies [7]. Tsetse flies of the *morsitans* group such as *G. pallidipes* have been shown to be the main vectors for the trypanosomes which cause Nagana in cattle, mainly: *T. b. brucei* and *T. congolense*[8]. Furthermore, the movement of cattle infected with *T. b. rhodesiense* has been implicated in new outbreaks of HAT in Uganda [9]. Africa Animal Trypanosomiasis is prevalent in some of the poorest areas in Sub-Saharan Africa and imposes a significant burden on development in this region [10].

Southern Nation Nationality and people Regional state has a total area of 438,370 hectare, which accounts for about 10 percent of the country. The main occupation of rural population is mixing farming practice where by crop and livestock are managed hand- in hand together. Cattle directly provide food such as meat, milk, by products such as hide. Indirectly they contribute over 30 percent to agricultural production by supplying essential input such as manure for replenishing soil fertility and restoring nutrient, and animal traction and power for ploughing and trashing; increasing the productivity of small holdings. Nevertheless, many factors affect the maximum benefit to be obtained cattle. Livestock disease is among the major factors that affect production and productivity, and trypanosomosis is the most important disease that influences livestock productivity in the region [11].

In 1970, *G. pallidipes* was recorded from lower Omo River and on the Wotto River and at Keiafer (1550 masl) also near Bako in GamoGofa province. It was found along the Segen River near Lake Chamo in GamoGofa. The lower Omo was also infested by this species up to down wards as far as Omorate. In addition, it found the upper part of the GalanaDulei valley (Wotto) with the Maze River valley

(Daramalo) [12]. The whole Omo belts which are indicated above are infested by *G. pallidipes*. Various types of Tsetse surveys undertaken in KindoKoyisha and Boloso Bombe Woreda of Wolaita Zone and Zima Waruma PA of Dawro Zone revealed the presence of *G. pallidipes* [13, 14].

G. pallidipes is one of the main vectors that transmit the sleeping sickness to humans and Nagana to animals [15]. It is therefore believed that *G. pallidipes* plays a central role in the disease burden of both HAT and AAT [16]. Current vector control interventions involve the use of insecticides either through sequential aerosol spraying technique (SAT); ground spraying; insecticide-treated targets or insecticide treated animals - live baits; the use of traps, and the sterile insect technique (SIT) [17].It was important that the knowledge of the daily activity and seasonal abundance of this particular species, *G. pallidipes* would be fundamental importance to determine what hour of the day and what month of the year to apply effective eradication strategy against it. The main objectives of this study were to investigate the monthly and daily activity pattern of *G. pallidipes* in study area to apply effective control and eradication measures.

Materials and Methods

Study Area description

The study was conducted from November 2018 to May 2019 in Maze National park at Gamo Zone, Southern Nations, Nationalities and Peoples' Regional State (SNNPRS), southwest Ethiopia. The Maze National park was located at 460 km and 235km south west of Addis Ababa and Hawassa, respectively in Gamo zone. The name of the park "Maze" is derived from the largest river that crosses the park. Maze River rises from southern parts of the surrounding highland and passes through the Maze national park from south to north direction and eventually drains in to Omo River. The study area Maze national park located $6^{\circ}25'$ N latitude and $37^{\circ}14'E$ longitude in Gamo zone and altitude ranges from 900 meter to 1400 meter above sea level [18].

The annual rainfall is between 843.8 mm and 1375.3 mm. The area experience a long rainy season that extends from April to October in which high amount of rainfall is recorded between April and June. The dry season is from November to march with mean maximum temperature $32.88^{\circ}C$. The cooler months in

the area include June, July and August with mean minimum temperature of 17.5 °C. Maze national park and its surroundings have unique natural, cultural, and historical attractions wide range of wildlife and vegetation types. The district should have different vegetation types were covered namely, woodland, savanna grassland, bushland, riverine forest and mixed types (woodland and grassland). The different wild animals found in the study area include: Baboon monkey, warthog, bush pig, antelope, bushbuck, lion, waterbuck, porcupine, greater kudu, lesser kudu, leopard, buffalo, wild cat other carnivores, and hippopotamus. Domestic animals in the study area include zebu cattle and goats [18].

Study population

The study populations for the study were tsetse flies (*G. pallidipes*) that occur in Maze National park at Gamo Zone, Southern Nations, Nationalities and Peoples' Regional State (SNNPRS), southwest Ethiopia.

Study Design

Longitudinal study was conducted to determine the seasonal and daily activity pattern of tsetse flies (*G. pallidipes*) in Maze national park. The study populations were collected based on sex, temperature, and relative humidity into account. Sample collection method was by using conical trap having different visual and odour attractants.

Study Methods

Monthly activity pattern of fly

For sampling *G. pallidipes*, six biconical traps [19] were used. The traps were set along transect from the open savanna grassland and riverine forest; sampling was taken every month for five consecutive days per month from November 2018 to May 2019. The distance between traps was 200 meter [20]. Each trap site was cleared of vegetation sufficiently to ensure reasonable visibility of the trap and dispersion of the odour attractants. Acetone plus cow urine was used as odour baits. The release rate of 500–600mg/h for acetone and 1–1.2g/h for cow urine were applied [21]. The trap has small cone that was placed on the upper part of the large cone where metal cage covered with white netting mashes was kept. The lower part is freely open for flies' entrance. The trap was held in place by the metal pole, which passes through the

wide opening. The lower part of the pole above the ground was greased to protect ants. To evaluate the environmental determinants of activity, hourly temperature and relative humidity were simultaneously recorded using temperature and moisture sensor thermo hygrometer.

Daily activity pattern of the fly


The area where tsetse population is high was selected for studying daily activity patterns. The study was done for three consecutive days over four month's period. For this, six biconical traps were set up and the distance between the traps was 100 meter. The number of tsetse flies caught in the traps was recorded every hour from dawn to dusk. Catch was made only for clear days, without rainfall and cloud, since the activity of tsetse flies are affected by weather condition of the area [22]. Trap catches were collected at hourly interval from 6:00am to 6:00 pm hour. To evaluate the environmental determinants of activity, hourly temperature and relative humidity were simultaneously recorded using temperature and moisture sensor thermo hygrometer. The flies caught were killed by gently squeezing their thorax then identified, sexed and counted.

Statistical Analysis

The data was analyzed using STATA statistical software program(window version 11). The seasonal fluctuation of *G. pallidipes* was analyzed using one-way analysis of variance (ANOVA). The daily activity pattern *G. pallidipes* was obtained by plotting the mean hourly catches \pm S.E. against hour of the day. Pearson correlation analysis was used to relate fly catch with temperature and relative humidity.

Results

Seasonal activity of *G.pallidipes* occurred with a monthly mean of 26.22 ± 1.59 SE. Its population built up during the rainy period (fig 1 and table 1). During the study period *G.pallidipes* population peaked during the rainy months April and May (Table 1). There was statistically significant variation in population during the five months ($F = 31.79$, $DF = 4$ and $p = 0.000$) (Table 2). There was also a positive correlation between fly abundance and rainfall during the rainy season.

Fig1. Monthly infestation of *G. pallidipes* in the maze national parkTable 1. Monthly variations in *Glossinapallidipes* apparent density during the study period in maze national park

M	o	n	t	h	R H %	mean	± S E	Tem.	°C	mean	± S E	Catch	mean	± S E
D	e	c	e	m	b	4 0 . 2 8	± 1 . 7 0	3 0 . 6 6	± . 8 4	2 1 . 1 9	± 1 . 0 4			
J	a	n	u	a	r	2 8 . 8 9	± 1 . 8 9	3 5 . 1 0	± 1 . 7	1 5 . 0 6	± 0 . 9 5			
M	a	r	c	h	3	3 1 . 3	± 1 . 3	3 6 . 4 8	± . 6 8	8 . 6 8	± . 4 6			
A	p	r	i	l	6	0 . 2 6	± 1 . 5	2 8 . 0 7	± . 8 3	4 0 . 6 0	± 1 . 6 9			
M	a	y			5	4 . 0 4	± 1 . 5	2 7 . 9 6	± . 8 1	4 4 . 3 3	± 1 . 7 2			

Table 2. Correlation of *G. pallidipes* apparent density with some ecological factors in study area.

Variables	M e a n	± S E	F	D	f	P -	v a l u e
Temperature	31.70	± 0.98	1 4	. 3 5 4	0	.	0 0 0 0 0
R H	45.1	± 1.58	9 1	. 3 1 4	0	.	0 0 0 0 0
C a t c h	26.22	± 1.59	3 1	. 7 9 4	0	.	0 0 0 0 0

The daily activity profile for *Glossina pallidipes* at maze national park shows two prominent peaks during the study period, a morning and afternoon peak. The activity increased gradually in the morning and peaked at 5:00 hour before declining between 6:00 and 10:00 hour local time. Their activity in evening was more vigorous, increasing steadily to peak at 12:00 hour and presumably continuing in the night.

Daily activity of *G. pallidipes* showed a bi-modal activity (Fig. 2). The relation between fly activities

and the prevailing temperature and relative humidity are shown in Table 3. There was a positive correlation of fly activity with temperature below 36°C. However, the relationship was insignificant ($r=0.699$, $p=0.122$). When the temperature exceeded 36°C there was negatively correlated ($r = -0.649$, $p = 0.023$) and the relationship was significant (Table 3). The relationship between fly activity and relative humidity was negatively correlated in the morning ($r = -0.649$, $P=0.1626$) and positively correlated in the afternoon ($r = 0.989$, $P = 0.002$).

Table 3. Correlation coefficients showing the relationship between fly activity with temperature and relative humidity in the morning (1-6) and afternoon (7-12)

Variables	Hour	Temp (°c)/RH%	Correlation coefficient	P – value
Temperature Vs catch	1 – 6	23.93 – 36.24	0 . . 6 9 9	0 . 1 2 2
Relative humidity Vs catch	1 – 6	53.27 – 29.93	- 0 , 6 4 9	0 . 1 6 2 6
Temperature Vs catch	7 - 12	33.44 – 45.76	- 0 . 8 6 7	0 . 0 2 5 6
Relative humidity Vs catch	7 - 12	22.76—34.34	0 . 9 8 9	0 . 0 0 0 2




Fig 2. Daily activity of *G. pallidipes* in maze national park

Discussion

From the present investigation, *G. pallidipes* were more abundant during the rainy months (April and May). Tsetse populations exhibit various patterns of population fluctuation related to local climates and vegetation. In general, pattern of fluctuations in tsetse populations is closely related to rainfall distribution throughout the year with a basic feature of an increase when rains start and decrease during the dry season [4]. In the present study, *G. pallidipes* populations increased at the start of the rains to reach a peak and then declined as the dry season starts.

The relatively low density of *G. pallidipes* during the dry months may be due to an accelerated consumption of its fat stored for further development of the pupa presumably due to high temperature associated with the low moisture content of the soil, which adversely affects the development of the puparium to the adult

stage [6]. The tsetse fly distribution is usually affected mostly by the rainfall and vegetation cover [23]. Similarly the present study shows that, during the wet season, especially during April to early May, flies were widely distributed when the vegetation cover of the area rehabilitate. This in turn provided shade and maintained a suitable microclimate for the flies and a better habitat for their vertebrate hosts. However, during the dry season, they have restricted distribution because of the scanty vegetation particularly shrubs.

The present study shows that *G. pallidipes* exhibited bimodal dry season activity for both sex. Unlike the case in the dry season, the activity profile during the wet season was unimodal and showed steady increases with time throughout the day. Since flies were compelled to concentrate in restricted areas where there is suitable habitat for their breeding and survival, attempt to control their population can be effective if conducted during dry season.

According to Dransfieldet al. [24] about 99% reduction can be obtained for *G. pallidipes*, with baited traps applied during the dry season. Rainfall is known to affect puparial development directly and the eco-distribution of the fly and the mammalian hosts indirectly [6]. The fact that the abundance of flies was statistically significant with rainfall (Table 2) indicates that the fly density increases as the rainfall increases.

Daily activity profile for *G. Pallidipes* at maze national park shows two prominent peaks during the study period (December to March), a morning and afternoon peak. The activity increased gradually in the morning and peaked at 4:00 hour before declining between 6:00 and 10:00 hour local time. In the evening, the flies were more vigorous, increasing steadily to peak at 12:00 hour and presumably continuing in to the night. Similarly, the activity profile for combined male and female *G. Pallidipes* at Nguruman- Kenya showed two prominent peaks during the dry season, while at wet season *G. pallidipes* did not show any prominent peak in the morning and afternoon [25]. In addition, Owaga et al.[26] observed that the activity of *G. pallidipes* at the Kenyan coast was throughout the day but with significant peak of activity between 9:00 and 10:00 and between 14:00 and 17: 00 hour. Most tsetse species are not active at night, but *G. pallidipes* was caught after 19.00 hour, i.e. 1:00 at the evening in local time [27]. In line with this, Fuller [28] also caught this species and *G. longiperinis* at night in Ethiopia when there were no lights; but the weather was very hot.

The result of the current study on the daily activity profile is also in accordance with other works [29-33]. Both sexes reached peak during the same hours of the day. Moreover, climatic factors influence the activity pattern to some extent [34]. Starvation also has an enormous effect on activity [35, 36]. Therefore, activity pattern may be unimodal or bimodal depending on whether optimal levels of abiotic and biotic factors have been exceeded or not, under particular circumstances.

There was a dual relation between the activity and temperature (Table 3). There was a positive correlation in the morning i.e., up to 6 am local time when the temperature was below 36°C and a negative correlation in the afternoon starting from 7:00 local time. This implies that the activity of *G. pallidipes* increased with a rise of temperature during the morning. This result agrees with that of Popham and

Vickers [37] who reported that tsetse flies are positively photo tactic below 30°C. The morning peak is followed by mid-day depression starting from 6 am up to 9 pm local time when the temperature was greater than 36°C. At higher temperatures the flies begin to seek shade [38]. During the afternoon, the fly number increased starting from 10:00 pm, when the temperature was less than 32°C. Like as tsetse flies have an innate pattern of activity. The savanna species, such as *G. morsitans* and *G. pallidipes*, are active mostly for the first two hour and the last two hours of the day [39]. These innate rhythms are influenced by temperature and thus the cool of morning in maze national park, when temperature less than 23°C can suppress the early morning peak of activity. Conversely, high (greater than 36°C) afternoon temperature, such as occur in maze national park can suppress the afternoon activity. But tsetse needs to feed and so when one peak was suppressed then the other compensates. For example, on very hot days in study area, tsetse flies are most active in the early morning.

Conclusion and Recommendations

From the investigation, it was clear that the reaction of *G. pallidipes* to abiotic and biotic factors is particularly important to their spatio-temporal distribution. The result showed that vector activity was majorly influenced by temperature and relative humidity and the activity pattern was sensitive to season. There was a seasonal change in habitat and abundance of the flies. They were more abundant during the rainy season and active around noon and in the evening. The main activity period is also dependent on temperature. Based on the conclusion, the following recommendations are forwarded:

- ❖ To improve agricultural development and livestock production, strategic vector control should be strengthened to eradicate vector depending on the activity pattern.
- ❖ It is also advisable to undertake intervention program during the dry season.
- ❖ Further study, particularly involving all the months, should be conducted in a wider area where the tsetse flies are known to cause significant impact on cattle production.

Acknowledgments

First of all, I would like to provide my glorious gratitude for the almighty GOD for his unlimited love, mercy and support up on me in every manner throughout my life. Next, my pleasure to Dr. Berhanu M. for his intellectual advice and devotion of his time to correct every mistake and I would thanks to Arba Minch tsetse flies and trypanosomiasis investigation and control center. Finally I would like to express my deepest gratitude to my families for their moral and financial supports.

References

1. Vreyen, M.,2001.Principles of area-wide integrated tsetse fly control using the sterile insect technique. Medicine Tropical., 61: 397-411.
2. Moloo, S. ,1993.The distribution of Glossina species in Africa and their natural hosts. Insect Science and its Application 4:511–527.
3. Abebe, G., J. B. Maloneand A .R. Thompson,2004. Geospatial forecast model for tsetse transmitted animal trypanosomosis in Ethiopia. SINET, Ethiopia. Journal. Science: 27 (1): 1-8.
4. Challier, A.,1982.The ecology of tsetse (Glossina spp.) (Diptera: Glossinidae). A review (1970– 1981). Insect Science and its Applications 3(2– 3):97–143.
5. Nagel, P. ,1995. Environmental Monitoring Hand Book for Tsetse Control Operations. MargrafVerlag, 323 pp.
6. FAO ,1982. Food and Agriculture Organization of the United Nations,(Pollock, J.N., ed.) Tsetse Control Training Manual, 1, Tsetse Biology, Systematic and Distribution, Techniques,Rome, 280.
7. Aksoy,S. andR.V.Rio,2008.Interactions among multiple genomes: tsetse, its symbionts and trypanosomes. Insect BiochemMol Biol. 2005 Jul;35 (7):691-8. doi: 10.1016/j.ibmb.2005.02.012. Epub 2005 Mar 28. Erratum in: Insect BiochemMol Biol. Nov;38 (11):1033. PMID: 15894186.
8. Geiger, A., F.Ponton andG. Simo,2015. Adult blood-feeding tsetse flies, trypanosomes, microbiota and the fluctuating environment in s ub Saharan Africa. The ISME Journal, 9(7): 1496–1507.
<http://doi.org/10.1038/ismej.2014.236>
9. Fevre,E., P.Coleman, M.Odiit, J.Magona, S.Welburn and M.Woolhouse,2001. The origins of a new Trypanosoma brucei rhodesiense sleeping sickness outbreak in eastern Uganda. Lancet, 358(9282): 625–628.
[http://doi.org/10.1016/S0140-6736\(01\)05778-6](http://doi.org/10.1016/S0140-6736(01)05778-6)
10. WHO,1998.Control and Surveillance of African Trypanosomiasis. WHO Technical Report Series 881. apps.who.int/iris/bitstream/10665/42087/1/WHO_TRS_881.pdf
11. CSA,2010): .central statistical agency, Federal democratic republic of Ethiopia agricultural sample survey. Bulletin, 446: 85-87.
12. Langridge, W.P. ,1976): .A Tsetse and Trypanosomosis Survey of Ethiopia. Ministry of Overseas Development Report, MOD, 100 pp.
13. Birhanu, A. ,1995. Preliminary survey on tsetse distribution and prevalence of bovine trypanosomosis in selected weredas of North Omo and KembataAlabaTambaro zones. AAU, Faculty of Veterinary Medicine, DebreZeit, DVM Thesis.
14. SRVL, (Southern Regional State Veterinary Laboratory),2004.1994-1998 Annual Reports. WolaitaSodo, Ethiopia.
15. Simarro, P., G.Cecchi, J.Franco, M.Paone, A.Diarra, J. Ruiz-Postigo and J.Jannin,2012.Estimating and Mapping the Population at Risk of Sleeping Sickness. PLoS Neglected Tropical Diseases, 6(10).
<http://doi.org/10.1371/journal.pntd.0001859>
16. Rogers, D. and T.Robinson,2004.Tsetse Distribution. In The Trypanosomes MA: Maudlin I, Holmes, P.H. & Miles (139– 179).
17. WHO,2011. Vector control Human African tryp anosomiasis. <http://www.who.int/mediacentre/factsheets/fs259/en/>.
18. Wegene, G. and W. Feleke,2015.land use practice, woody plant species diversity and associated impacts in maze national park, southeast Ethiopia. Plant. 3: 64-74.

19. Laveissier, C. and P. Grebaut,1990).: Recherchesurlespieges a Glassines (Diptera: Glossinidae) miseau point d'un mode'leeconomique le piege 'Vavoua'. Tropical Medicine and Parasitology41:185–192.
20. Steve, M. ,1997. Practical trapping guidelines. International Centre of Insect Physiology and Ecology, Nairobi, Kenya.In: Tsetse and Trypanosomosis Literature Notes
21. Vale, G., D. Halland T. Gough,1988.The olfactory response of tsetse flies, spp. (Diptera: Glossinidae) to phenols and urine in the field. Bulletin of Glossina Entomological Research78:293– 300.
22. Mohammed-Ahmed, M. and A. Odulaja,1997.Diel activity patterns and host preferences of *Glossinafuscipes* (Diptera: Glossinidae) along the shores of Lake Victoria, Kenya. Bulletin of Entomological Research 87:179–186.
23. Merid, N., G.Melaku and S.Emiru,2004.seasonal and daily activity pattern of *G. morsitanssubmorsitans*(Diptera: Glossinidae) in Ghibe river valley. SINET: Ethiopia journal Science.27(2): 135-142.
24. Dransfield, R., R.Brightwell, C. Kyorkuand B. Williams,1990.Control of tsetse fly (Diptera: Glossinidae) populations using traps at Nguruman, South-west Kenya. Bulletin of Entomological Research80:265–276.
25. Okoth. S., E.Kokwaro, J. Kiragu and G.Murilla,2007. Diurnal activity of allopatric population of *Glossinapallidipes* and host and their implication on risk of transmission of sleeping sickness in Kenya. Journal of entomology, 4(1): 20-32.
26. Owaga, L., Okello and M. Chaudhury,1993.Diel activity pattern of *Glossinaausteni* new stead (Diptera: Glossinidae) in the field and in the laboratory. Insect science. applied, 14: 701-705.
27. KangwagyeT. ,1971. Observations on *Glossinafuscipes*Newstead, *G. fuscipleuris* Austen and *G. pallidipes* Austen in Western Uganda. Organization of African Unity/Scientific and Technical Research CommissionIn Internatiortal Scientific Council for Trypanosomiasis Research.105: 187-191
28. Fuller G. ,1978. Distribution of *Glossina* (Diptera: Glossinidae) in south-western Ethiopia. Bulletin of Entomological Research. 68: 299-305.
29. Mulligan, H.,1970. The African Trypanosomosis. Ministry of Overseas Development and George Allen and Unwin, London, 950 pp.
30. Brady, J. ,1973.The pattern of spontaneous activity in tsetse fly *Glossinamorsitans*Westw. (Diptera: Glossinidae). Bulletin of Entomological Research 63:441–444.
31. Tesfa-Michael T. ,1980. Preliminary ecological studies of *Glossinamorsitans* (Glossinidae: Muscidae) in Didessa valley, Ethiopia. Ethiopian Journal of Agricultural Science2:129–138.
32. Turner, D.A.,1980. Tsetse ecological studies in Nigeria and Mozambique. Resting behaviour. Insect Science and its Application1:15–21.
33. GetachewT. ,1983. Studies on tsetse flies of Finchaa River Valley. MSc Thesis, School of Graduate Studies, Addis Ababa University, 83 pp.
34. Brady, J. and A. Crump,1978.The control of circadian activity rhythms in tsetse flies: environmental or physiological clock. Physiological Entomology3:177–190.
35. Barrass, R.,1970.The flight activity and settling behaviour of *Glossinamorsitans*Westw. (Diptera: Muscidae) in laboratory experiments. Bulletin of Entomological Research 59:627–635.
36. Crump, A.J. and J. Brady,1979. Circadian activity patterns in three species of tsetse fly: *Glossinapalpalis*, *G. austeni* and *G. morsitans*. Physiological Entomology4:311–318.
37. Popham, E.J. and H. Vickers,1979. Tsetse fly reactions to light and humidity gradients. Experientia35:194–196.
38. Pilson, R. and B.Pilson,1967.Behavioural studies of *Glossina morsitans*Westw. In the field. Bulletin of Entomological Research57:227–252.

39. Hargrove, J. and j. Brady, 1992. Activity rhythms of tsetse flies (*Glossina* spp.) (Diptera: Glossinidae) at low and high temperature in nature. Bulletin of entomological research, 82, 321-326.

Access this Article in Online	
	Website: www.ijarbs.com
	Subject: Entomology
Quick Response Code	DOI: 10.22192/ijarbs.2021.08.04.008

How to cite this article:

Abenezer Mara Zuma. (2021). Monthly and Daily Activity Pattern of Tsetse Fly *Glossina pallidipes* in Maze National Park, Gamo Zone, Southwest Ethiopia. Int. J. Adv. Res. Biol. Sci. 8(4): 55-63.
DOI: [http://dx.doi.org/10.22192/ijarbs.2021.08.04.008](https://doi.org/10.22192/ijarbs.2021.08.04.008)