



## **Hematological and Biochemical Investigations in Albino Rat Exposed to Cadmium Chloride.**

**Kshama Dwivedi**

Department of Zoology, Bareilly College, Bareilly - 243005(U.P.), INDIA

Corresponding Author. E-mail: [kshamadwivedi74@gmail.com](mailto:kshamadwivedi74@gmail.com)

### **Abstract**

Cadmium being a non-essential element is present as a contaminant in food, water, and polluted air. Concern about its possible effect on the general population exposed to low concentration over long periods of time has been raised in part because of its steadily increasing consumption and release in the environment. The present study was to investigate changes in hematological and biochemical parameters of albino rats exposed to cadmium chloride.

Animals were exposed with low and high doses of cadmium chloride, along control. They were treated for 30 and 60 days separately, in two experimental sets. In each of the experimental sets, Group A was control, Groups B exposed with lower dose of CdCl<sub>2</sub> (2.6mg/kg.b.wt.) and group C with high dose of CdCl<sub>2</sub> (5.2mg/kg.b.wt.). The findings of this study indicate that rats intoxicated with cadmium chloride for 30 days showed decrease in TEC, TLC, Hb and PCV (P<0.05) values and increase in MCV and ESR (P<0.05) values. Rats exposed for 60 days showed significant fall (P<0.05) in all these parameters except ESR. There was significant increase (P<0.05) in the values of serum creatinine, urea, aspartate aminotransferase (AST) and alanine aminotransferase (ALT) while decrease in the values of alkaline phosphatase (ALP), cholesterol, and protein in the blood samples of animals exposed to different doses of cadmium chloride as compared to control when treated for 30 and 60 days.

**Keywords:** Cadmium chloride, haematological parameters, biochemical parameters, liver, kidney, and enzymes.

### **Introduction**

Cadmium, a heavy metal is highly persistent environmental toxicant. It is present in air, drinking water as well as food and has potential to affect the health of human population; mainly those who live in highly industrialized regions (Sharrett *et.al.*, 1983, Smigiel *et. al.*, 1994). Cadmium is considered as an occupational and environmental pollutant. As food chain contaminant, toxicity of cadmium is of great concern (MacLaughlin *et. al.* 1999; ATSDR, USA, 2012). Cadmium is present in cigarette smoke and polluted air, contributing to elevated cadmium concentration in blood urine and tissues of smokers,

compared with non-smokers of similar age and gender (Satarug *et. al.* 2017)

Cadmium enters the body by various routes and gets accumulated in various organs and organ systems such as brain, liver, lungs, blood system, bone, spleen, pancreas, and testes (Cai *et. al.*, 2001; Emmanuel *et. al.*, 2003). Cadmium toxicity is reported to be nephrotoxic (Jarup *et. al.*, 2009), associated with generalized osteoporosis, osteomalacia (Horiguchi *et. al.*, 2010) and type-2 diabetes (Satarug *et. al.*, 2017).

Cadmium also has toxicological impact on plants and so their products for human consumption (Amir *et. al.*, 2014). Intensive agricultural practices and industrial processing has resulted in the contamination of food, feed, and water, becoming source of exposure for humans. Some geographical areas are associated with the naturally occurring high cadmium concentration. Historically, rice consumption which being contaminated with cadmium from zinc discharge caused an outbreak of itai-itai disease affecting mostly women in Japan (Baba *et. al.*, 2014).

Food and Agricultural organization (FAO) and World Health Organization (WHO) joint expert committee established guideline of a safe dietary cadmium intake and a urinary cadmium threshold limit. Currently dietary cadmium intake is within the FAO/WHO tolerable level of 58µg/day for 70 kg person (Soisungwan Satarug, 2018). But industrial pollution, other source of cadmium exposure like cigarette smoke, ingestion of polluted vegetables and ambient air in urban- industrialized areas (Gallagher *et. al.*, 2004 and Thomas *et. al.*, 2009) create a serious chronic exposure to heavy metals especially to cadmium. Asian subpopulations have been found to have the highest mean blood cadmium among five ethnic group studied in the U.S. National Health and Nutrition Examination Survey (NHANES 2011-2012). The aim of this study was to investigate the toxic effects of chronic and sub chronic cadmium exposure on hematological and biochemical parameters in albino rats. Experimental results will act as biomarker in the cadmium toxicity.

## Materials and Methods

### Experimental Animals

Albino rats of 6 to 10 weeks old weighing 150- 160 grams were purchased from the Laboratory Animal Resource Section, of Indian Veterinary Research Institute (IVRI) Izzat Nagar Bareilly, U.P. that were maintained in experimental animal shed of the division. Animals brought into new conditions were acclimatized to the new environment prior to the experiments. Rats were kept under conventional condition (6 rats per steel cage, 12 hr. light to dark cycle) and were maintained with standard rat food and tap water *ad libitum*. All the chemicals used were from the Sigma Chemicals Co., Merk and Qualigens.

### Experimental Design

The experimental rats were randomly divided into three groups A, B, and C each comprising of 6 animals. Group A was control remaining two groups were intoxicated with different chronic doses of heavy metal (Cadmium chloride) compound, Group B Cadmium chloride (low dose) with 2.6mg/kg.b.wt, Group C with Cadmium chloride (high dose) 5.2mg/kg.b.wt The compounds were given in tap water as they were easily soluble in it while the control received only plain tap water per os by gavage. Rats under the above treatment were monitored for 30 days and 60 days and their body weight were taken weekly. Mortality rate, food consumption, clinical signs and symptoms and behavioral activities were under observation.

Blood was collected for hematological and biochemical examination after 30 days and 60 days from the retro-orbital plexus with the help of capillary tube as described by Sorge and Buckner (1964). Blood was collected in two aliquots. In one aliquot, EDTA (1mg/ml) was added for hematological parameters estimation other aliquot was without any anticoagulant for harvesting serum for biochemical estimation (centrifuged at 3000 rpm for ten minutes). The test samples were stored in air vials at -20°C till used.

### Statistical analysis

All data presented as the mean  $\pm$ standard error of mean (SEM). The results were analyzed for statistical significance by one-way analysis of variance (ANOVA) followed by Dunnett's *post hoc* test of significance. *P* values less than 0.05 (*p* 0.05) were considered as statistically significant.

### Results and Discussion

Clinical observations showed that exposed animals were docile and less active than control group. No mortality occurred in control and other groups treated with the different doses of heavy metal. There was time and dose-dependent reduction in the body weight when treated for 30- and 60-days duration. The effect of treatment of cadmium chloride on different organs of albino rats when intoxicated for 30 days and 60 days is given in Table 1 to 6.

**Table 1:** Hematological profile of albino rats after oral administration (gavage) of cadmium chloride for 30 days.

Group	TEC (Millions/mm <sup>3</sup> )	TLC (Thousands/mm <sup>3</sup> )	Hb(g/dl)	PCV (%)	MCV(fL)	ESR (Mm <sup>3</sup> /L)
<b>Group-A:</b> Control	7.32±0.88	7.33±0.33	13.50±0.39	48.21±2.07	65.86±12.44	1.25±0.25
<b>Group-B:</b> Cd (L)	6.65±0.76	7.05±0.15	11.34±0.24	40.50±3.08	60.90±9.66	1.00±0.00
<b>Group-C:</b> Cd (H)	6.49±0.54	6.49±0.52	11.11±0.11	42.40±1.02	65.33±7.43	1.40±0.37

- Cd (L) = Cadmium Chloride Low Dose
- Cd (H) = Cadmium Chloride High Dose
- All the values are mean±SE; n=6

After 30 days post treatment, TEC showed trend of decrease. The maximum decrease after 30 days treatment period was observed in group C (11.34%). Group B showed minimum decline value of 9.15% as compared to the control. TLC values decreased to the maximum in group C (11.45%) and a significant fall

in PCV was observed having a value in group B (15.99%). The animals of group B showed significant fall in MCV of 7.53% and hemoglobin group C (21.62%) on contrary ESR increased insignificantly in both the treated groups.

**Table 2:** Differential leucocyte count (DLC) after oral administration (gavage) of cadmium chloride for 30 days.

Groups	DLC				
	Lymphocytes%	Neutrophils%	Monocytes%	Eosinophils%	Basophils%
<b>Group-A:</b> Control	70.11±2.14	21.66±2.65	5.50±0.67	1.33±0.30	0.92±0.22
<b>Group-B:</b> Cd (L)	67.15±3.21	23.24±0.13	7.15±0.33	1.74±0.21	0.97±0.17
<b>Group-C:</b> Cd (H)	66.36±0.40	23.89±4.03	6.97±0.33	1.83±0.31	1.14±0.14

- Cd (L) = Cadmium Chloride Low Dose
- Cd (H) = Cadmium Chloride High Dose
- All the values are mean±SE; n=6

Post treatment intoxication after 30 days showed fall in the lymphocyte number. Contrary to lymphocytes,

neutrophils, monocytes, eosinophils, and basophils shows increase in number compared to control.

**Table 3:** Hematological profile of albino rats after oral administration (gavage) of cadmium chloride for 60 days.

Group	TEC (Millions/mm <sup>3</sup> )	TLC (Thousands/mm <sup>3</sup> )	Hb (g/dl)	PCV (%)	MCV (fL)	ESR (Mm <sup>3</sup> /L)
<b>Group-A:</b> Control	7.59±0.13 <sup>a</sup>	7.29±0.08 <sup>a</sup>	13.57±0.24 <sup>a</sup>	48.63±0.62 <sup>a</sup>	64.07±0.40 <sup>a</sup>	1.18±0.03 <sup>b</sup>
<b>C.V.</b>	4.13	2.61	4.41	3.12	1.51	6.36
<b>Group-B:</b> Cd (L)	6.55±0.12 <sup>b</sup> <sup>c</sup>	6.38±0.12 <sup>bc</sup>	11.34±0.29 <sup>bc</sup>	39.52±0.60 <sup>b</sup>	60.33±1.32 <sup>b</sup>	1.09±0.03 <sup>bd</sup>
<b>C.V.</b>	4.78	4.62	6.47	3.71	5.37	7.60
<b>Group-C:</b> Cd (H)	6.15±0.18 <sup>b</sup>	6.24±0.19 <sup>b</sup>	10.62±0.21 <sup>bdf</sup>	37.13±0.87 <sup>b</sup>	59.33±0.62 <sup>b</sup>	1.27±0.04 <sup>ac</sup>
<b>C.V.</b>	7.18	7.29	4.93	5.75	5.58	6.95

- Cd (L) = Cadmium Chloride Low Dose
- Cd (H) = Cadmium Chloride High Dose
- All the values are mean±SE; n=6

At the end of 60 days treatment there was decrease of TEC in both the groups which was being intoxicated but in group C, with high dose of cadmium it was significant (18.9%). TLC lowered when compared to control. In group C and group B fall of 14.32% and 12.50% in TLC was observed. PCV and MCV also decreased but very significantly in group B and C. Fall of 24.7% in PCV was noticed in group C. Hemoglobin

percentage decrease in the animals of both the intoxicated groups compared to control and it was 16.3% in group B and 21.62% in group C. ESR on contrary increased in both the treated groups. In DLC leucocyte count lymphocytes decreased but significant increase was observed in eosinophils and basophils in both the 60 days intoxicated groups.

**Table 4:** Differential leucocyte count (DLC) after oral administration (gavage) of cadmium chloride for 60 days.

Groups	DLC				
	Lymphocytes%	Neutrophils%	Monocytes%	Eosinophils%	Basophils%
<b>Group-A:</b> Control	70.91±1.10	21.50±0.94	5.50±0.11 <sup>bdfhj</sup>	1.17±0.13	0.92±0.04
<b>C.V.</b>	3.79	10.71	4.78	26.23	10.69
<b>Group-B:</b> Cd (L)	63.64±1.55	26.10±1.64	7.38±0.10 <sup>bdfg</sup>	1.62±0.13	1.03±0.04
<b>C.V.</b>	5.98	15.39	3.56	18.95	9.55
<b>Group-C:</b> Cd (H)	61.04±1.09	27.36±1.92	8.58±0.28 <sup>c</sup>	1.76±0.09	1.17±0.04
<b>C.V.</b>	4.41	17.19	8.20	13.04	8.41

●Cd (L) = Cadmium Chloride Low Dose

●Cd (H) = Cadmium Chloride High Dose

●All the values are mean±SE; n=6

When blood biochemical parameters were estimated the animals receiving cadmium chloride showed significant increase in the values of AST, ALT, creatinine, and urea when treated for 30 days. Group C increase showed more significant increases in AST

(6.07%), ALT (28.82%), creatinine (30.76%) and urea (18.37%) as compared to control. On the other hand, ALP, cholesterol, and protein values showed marked decrease from 16.9% to 30.8% in these parameters. It was strange to see fall in ALP.

**Table 5:** Blood biochemical profile after oral administration (gavage) of cadmium chloride for 30 days.

Group	AST (IU/L)	ALT (IU/L)	ALP (IU/L)	Creatinine (mg/dl)	Cholesterol (mg/dl)	Urea (mg/dl)	Protein (g/dl)
<b>Group-A:</b> Control	39.5±1.21	28.1±0.28	57.00±0.40	0.65±3.53	20.22±1.52	29.6±1.6	7.44±0.19
<b>Group-B:</b> Cd (L)	40.4±0.96	31.2±0.91	46.02±0.72	0.67±1.15	17.08±2.69	31.9±1.51	6.50±0.62
<b>Group-C:</b> Cd (H)	41.9±2.03	36.2±2.15	39.41±0.40	0.85±1.06	16.46±3.27	35.04±2.1	6.18±0.52

●Cd (L) = Cadmium Chloride Low Dose

●Cd (H) = Cadmium Chloride High Dose

●All the values are mean±SE; n=6

**Table 6:** Blood biochemical profile after oral administration (gavage) of cadmium chloride for 60 days.

Group	AST (IU/L)	ALT (IU/L)	ALP (IU/L)	Creatinine (mg/dl)	Cholesterol (mg/dl)	Urea (mg/dl)	Protein (g/dl)
Group-A: Control	40.1±1.46	30.1±0.24	54.92±1.50 <sup>a</sup>	0.71±1.08	20.95±0.44	30.43±2.38	6.60±0.11
Group-B: Cd (L)	41.87±1.45	32.7±2.59	39.98±1.35 <sup>be</sup>	0.89±2.31	18.69±1.58	33.75±1.86	6.28±0.33
Group-C: Cd (H)	42.4±2.5	41.5±2.3	30.33±1.71 <sup>bdf</sup>	1.00±1.36	15.87±1.44	41.77±1.41	5.90±0.31

●Cd (L) = Cadmium Chloride Low Dose

●Cd (H) = Cadmium Chloride High Dose

●All the values are mean±SE; n

In the second set of experiment for 60 days the AST, ALT, creatinine, and urea values showed the same trend of increase in cadmium intoxicated groups i.e., Group B and Group C. Animals of group C with high dose of cadmium showed increase of 5.7% in AST level, 37.8% in ALT, 40.84% in creatinine and 37.26% in urea when compared to control group A. Cholesterol, ALP, and protein showed decrease in both the treated group compared to control. Fall of 44.77% in ALP, 24.44% in cholesterol and 10.6% in protein was observed. Group B showed less fall when compared to group C with high cadmium dose. Decrease observed was dose and duration dependent. These observations shows that cadmium produced toxic effect on body hampering the function of liver kidney and blood parameters.

Clinical observations showed that animals were docile and less active than control. No mortality occurred in control as well as intoxicated groups with cadmium chloride. In treated group there was a dose-dependent reduction in body weight gain. Hematological and biochemical indices have been reported to be reliable parameters for the assessment of health status of animals and humans (Ohaeri *et. al.*, 2011). Toxic compounds affect the metabolism and function of mature blood cells and disturb hematopoiesis process. The response to toxic influences manifests itself primarily in reduction of the number of circulating blood cells along its functional and structural abnormalities.

Alteration in hematological parameters and activity of serum enzymes were frequently indicators of toxicity, organ damage and cell damage (Kodavanti and Mehendale, 1991). Production of cells in bone marrow is reported to be highly sensitive to toxic influences.

Sometimes only one cell line is affected by toxic substances and sometimes all of them.

In the present study TEC, TLC, hemoglobin percentage, PCV and MCV declined sharply in all the groups when compared with control, but ESR significantly ( $p < 0.05$ ) increased in cadmium intoxicated rats. These were in accordance with Guilhermino *et. al.* (1998) and Hounkaptin *et. al.* 2013. Fall in TEC was also reported by Guedenon *et. al.* (2012) on the fishes exposed to cadmium and mercury. Low values of TLC in treated groups are related to toxic action of cadmium and mercury which can induce leukopenia and thrombocytopenia in cases of severe liver dysfunction (Lee *et.al.*, 2004) in mice treated with lead (Lodia and Kansala, 2012; Veena *et. al.*, 2011). Being highly sensitive to toxic influences hematological system results in abnormal production or inhibition of production of blood cells (Marx, 1996; ATSDR, 1999). The observed decrease in erythrocytes, hemoglobin and hematocrit is consistent with the previous studies by Horiguchi *et. al.* (2007). Anemia might have caused not only by decrease in the synthesis and release of erythrocytes in the blood circulation (Vinodhini and Narayanan, 2008) but also by destruction of erythrocytes (Kori-Siakpere, 2009). Hounkpatin *et. al.*(2013) and Wakeel *et al.*(2020) also reported decrease in RBC, MCV, PCV and hematocrit in cadmium exposure of rats.

Many biochemical parameters are biomarkers to evaluate the possible toxic effect on liver and kidney caused due to disturbed physiological functions responsible for many diseases (Smaoui *et. al.*, 2000; Hannah *et. al.*, 2016). The result of study showed a significant increase in enzymatic activity of AST and ALT.

Increased activity of these enzymes in blood represents their leakage of tissue into plasma following a hepatic lesion responsible for the deterioration of the membrane permeability (Layachi and Kechrid, 2012; Diaby *et. al.*, 2016) and characteristic white globules (Albasha and Azab, 2014) destroying cell of liver directly. It is also reported to be cellular lesion and pyknotic nuclei caused due to the cell necrosis in cadmium toxicity (Wakeel *et. al.*, 2020). Results of this study shows significant increase in the level of creatinine and urea compared to the control, may be due to renal insufficiency (Guilhermino *et. al.*, 1998; Boujelbene *et. al.*, 2002). Kidneys are reported to be target organs of toxic effects of cadmium (Wang *et. al.*, 2014; Satarug, 2018) which accumulate here causing kidney damage (Poontawee *et. al.*, 2016). Cadmium toxicity causes tubular dysfunction reducing glomerular filtration leading to kidney failure (Fahim *et. al.*, 2012). Disrupted nuclei, paler cytoplasm, and scanty chromatin (Wakeel *et. al.*, 2020).

Cholesterol, ALP, and plasma protein decreased in the treated animals was also reported by Guilhermino *et. al.* in 1998. Fall in ALP as biomarker might be associated to hepatobiliary injury. The decrease in plasma total protein, which seems to be attributed to decrease in albumin was also indicative of cadmium induced liver and kidney injury (Guilhermino *et. al.*, 1998). Ferrao *et. al.* (2010) reported that blood cadmium is associated to impaired renal function and proteinuria (albuminuria). In United States population, low cholesterol is associated with the lipid metabolic abnormality due to the cadmium toxicity agrees to the study by Saeed Samarghandian *et. al.* (2015), who reported that even low cadmium concentrations can adversely, affects the lipid and lipoprotein profile is a lipidperoxidation. Zhou *et. al.* (2016) reported that cadmium exposure is associated with dyslipidemia.

## Conclusion

The results obtained from the present study indicates that cadmium toxicity in chronic and sub chronic dose induce alteration in different blood parameters. Cadmium affects the blood by lowering cell count and volume leading to anemia. Increase in AST, ALT, creatinine, and urea showed that cadmium produces toxic effects on liver and kidney. Low cholesterol,

ALP, and blood protein was observed. Thus, the study shows that cadmium as environmental pollutant is hazardous to human and animal health. Monitoring the level of these heavy metal contaminant in eatables, drinking water and occupational exposure, humans can be saved from disease and ill health leading to sustainable environment.

## Acknowledgments

The author is thankful to Dr. V. P. Varshney, Principal Scientist (NRL Division), Dr. H. C. Tripathi, Principal Scientist (Pharmacology and Toxicology division) and Dr. D. K. Gupta, Associate Professor, Department of Zoology, Bareilly College, Bareilly for providing facility to carry out this work and cooperation.

## References

1. Albasha, MO. And El-Saied Azab, A. (2014). Effect of cadmium on liver and amelioration by aqueous extracts of fenugreek seeds, rosemary, and cinnamon in guinea pigs: Histological and biochemical study. *Cell biology* 2(2): 7-17.
2. Amir, W., Jahanzaib, A., Farhat, I., Ashif, S., Zahid, M., and Gulam, M. (2014). Pollution status in Pakistan: A retrospective review on heavy metal contamination of water, soil, and vegetables. *Biomedical Research international*. 29 pages.
3. ATSDR (Agency for Toxic Substances and Disease Registry).(1999). ATSDR HazDat database. Atlanta, G. A: US. Department of Health and Human Services.
4. ATSDR (Agency for Toxic Substances and Disease Registry).(2012). Toxicological Profile for Cadmium; Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention: Atlanta, GA, USA.
5. Awata, H., Linder, S., Mitchell, L. E., and Delclos, G. L. (2017). Biomarker levels of toxic metals among Asian population in the United States: NHANES 2011-2012. *Environmental Health Perspective*.125: 306-313.
6. Baba, H., Tsuneyama, K., Kumada, T., Aoshima, K., and Imura, J. (2014). Histological analysis for osteomalacia and tubulopathynin itai-itai disease. *Journal of Toxicological Sciences*. 39: 91-96.

7. Boujelbene, M., Gorbel, F., Ayadi, F., Makni-Ayadi, F., Makni-Guermazi, F., Kammoun, A., Croute, F., Soleilhavoup, J. P. and Eleki, A. (2002). *Impact d'un*.
8. Cai, Y., Aoshima, K., Katoh, T., Teranishi, H. and Kasuya, M. (2001) Renal tubular dysfunction in male inhabitants of cadmium-polluted area in Toyama, Japan, eleven-year follow-up study. *Journal of Epidemiology*. 11:180-9.
9. Diaby, V., Yapo, A. F., Adon, A. M., Dosso, M. and Djama, A. J. (2016). Renal, hepatic, and splenic biotoxicity of cadmium sulphate in wistar rats. *International Journal of environmental Science and Toxicology Research*. July,4(6): 103-110.
10. Emmanuel, E., Allison, K., Margaret, M., Carmen, A., Elizabeth, A., Kanessa, L. W., and Bhattacharyya, M. H. (2003). Cadmium pathways during gestation and lactation in control versus metallothionein 1,2- knockout mice. *Toxicol Sci*. Feb;71(2):154-63.
11. Fahim, M. A., Nemmar, A., Dhanasekaran, S., Singh, S., Shafiullah, Y. M., Zia, S. and Hasan, M. Y. (2012). Acute cadmium exposure causes systemic and thromboembolic events in mice. *Physiological Research* 61: 73-80.
12. Ferraro, P. N., Costanzi, S., Naticchia, A., Sturniolo, A. and Gambaro, G. (2010). Low level exposure to cadmium increases the risk of chronic kidney disease: analysis oh the NHANES 1999-2006. *BMC Public Health*. 10; 304:1-8.
13. Gallagher, C. M., Kovach, J. S., and Meliker, J. R. (2004). Urinary cadmium and osteoporosis in U. S. women >or =50 years of age: NHANES1988-1994 and 1999-2004. *Environmental Health Perspective*.116: 1338-43.
14. Guedenon, P., Etorh, P. A., Hounkpatin, A. S. Y., Alimba, C. G., Ogunkanmi, A., and Nwokejegbe, E.C. et al. (2012). Haematological study of *Clarias gariepinus* exposed to chronic and sub chronic doses of cadmium, mercury and combined cadmium and mercury. *Sci. Nat*. 4(2);2-19
15. Guilhermino, L., Soares, A. M. V. M., Carvalho, A. P., and Lopes, M. C. (1998). Effect of cadmium and parathion exposure on hematology and blood biochemistry of adult male rats. *Bull. Environ. Contam. Toxicol*. 60: 52-59.
16. Hannah, W.N., Torres, D. M., and Harrison, S. A. (2016). Nonalcoholic steatohepatitis and endpoints in clinical trials. *Gastroenterology and Hepatology*. 12(12): 755- 763.
17. Horiguchi, H. (2007). Anemia induced by cadmium intoxication, *Nihon Eiseigaku Zasshi. Japanese Journal of Hygien*. 62(3): 888-904.
18. Horiguchi, H., Aoshima, K., Oguma, E., Sasaki, S., Miyamoto, K., Hosoi, Y., Katoh, T. and Kayama, F. (2010). Latest status of cadmium accumulation and its effects on kidneys, bone and erythropoiesis in inhabitants of the formerly cadmium polluted Jinzu river basin in Toyama, Japan, after restoration of rice paddies. *Int. Arch. Occup. Environ. Health*. 83:953-970.
19. Hounkpatin, A. S., Etorh, P. A., Guédénon, P., Alimba, C.G., Ogunkanmi, A., Dougnon, T. V., Boni, G., Assi, K. A., Montcho, S., Loko, F., Ouazzani, N., Mandi, L., Boko, M. and Creppy, E. E. (2013). Hematological evaluation of Wistar rats exposed to chronic doses of cadmium, mercury and combined cadmium and mercury. *African Journal of Biotechnology*.12(23): 3731-3737.
20. Jarup, L., and Akesson, A. (2009). Current status of cadmium as an environmental health problem. *Toxicology and Applied Pharmacology*. 238; (3): 201-8.
21. Kodavanti, P. R. and Mehendale, H. M. (1991). Biochemical methods of studying hepatotoxicity. Meeks, R. G., harisson, S. D., Bull, R. J. In: *Hepatotoxicity* (Eds). CRC press, Boca Raton.
22. Kori-Siakpere, O., Ogbe, M. G. and Ikomi, R. B. (2009). Hematological response of the African catfish *Clarias gariepinus* (Burchell, 1822) to sublethal concentrations of potassium permanganate. *Sci. Res. Essay. Academic Journal*. 4 (5): 457-466.

23. Layachi, N. and Kechrid, Z. (2012). Combined protective effect of vitamin C and E on cadmium induced oxidative liver injury in rats. *African Journal of Biotechnology*. 11(93): 16013-16020.
24. Lee, W. M. (2004). Acetaminophen and the US acute liver failure study group: Lowering the risks of hepatic failure. *Hepatology*. 40: 6-9.
25. Lodia, S. and Kansala, L. (2012). Antioxidant activity of *Rubia cordifolia* against lead toxicity. *International Journal of Pharmacology Science Research*. 3(7): 2224-2232.
26. Marx, J. J. M. (1996). Toxicology of the blood, pathophysiology, toxicological pathology and mechanistic aspects. Niesink R. J. M., Vries J. D., Hollinger M. A. In: *Toxicology Principles and Applications*. Eds Raymond.
27. McLaughlin, M. J. and Singh, B. R. (1999) Cadmium in soils and plants. In *Developments in Plants and Soil Sciences*; McLaughlin M. J., Singh B. R., Eds; Kluwer Academic Publisher: Dordrecht, The Netherlands; Boston, London: Volume 85: 1-7.
28. Ohaeri, C. C. and Eluwa, M. C. (2011). Abnormal biochemical and hematological indices in Trypanosomiasis as a threat to herd production. *Veterinary Parasitology*. 177: 199-202.
29. Poontawee, W., Surapol, N., and Orawan, W. (2016). Protective effect of *Cleistocalyx nervosum* var. *paniala* fruit extract against oxidative renal damage caused by cadmium. *Molecules*. 21(133): 1-13.
30. Samarganghian, S., Nezhad, M. M. S., Azad, F. J., Farkhondeh, T. and Bafandeh, F. (2015). Effect of chronic exposure to cadmium on serum lipid, lipoprotein and oxidative stress indices in male rats. *Interdisciplinary Toxicology*. Sep; 8(3): 151-154.
31. Satarug, S., Vesey, D. A. and Gobe, G. C. (2017). Current health risk assessment practice for dietary cadmium: Data from different countries. *Food and Chemical Toxicology*. 106: 430-445.
32. Satarug, S. (2018). Dietary cadmium intake and its effects on kidneys. *Toxics*. 6 (15): 1-23.
33. Sharrett, A. R., Carter, A.P., Orheim, R. M. and Fenlein, M. (1982). Daily intake of lead, cadmium, copper, and zinc from drinking water: The Seattle study of trace metal exposure. *Environmental Research*. Aug;(2):28,456-475.
34. Smaoui, M., Fatma, G., Manel, B. Fatma, M. A., and Abdel Fattah, E. (2000). Impact de l'exposition chronique aux gaz d'échappement d'origine automobile sur certains biomarqueurs touchant la fonction hormonale sexuelle mâle, la fonction rénale at l'hémogramme chez le rat. *Pollution Atmosphérique*. 167: 439-449.
35. Smigiel, D. (1994). Accumulation of heavy metals (Pb, Cd) in selected varieties of vegetables. *Rocz. Panstw. Zakl. Hig.* 45: 279-284.
36. Sorg, D. A. and Buckner, B. (1964). A simple method of obtaining venous blood from small laboratory animals. *Proc. Soc. Exp. Bio. Med.* 115: 1131-1132.
37. Thomas, L. D., Hodgson, S., Nieuwenhuijsen, M., and Jarup, L. (2009). Early kidney damage in a population exposed to cadmium and other heavy metals. *Environmental Health Perspective*. 117;181-4.
38. Veena, S., Leena, K., Arti, S., Shweta, L., and Sharma, S. H. (2011). Ameliorating effect of *Coriandrum sativum* extracts on hematological and immunological variables in an animal model of lead intoxication. *Journal of Pharmacology and Allied Health Sciences*. 1: 16-29.
39. Vinodini, R. and Narayanan, M. (2008). Bioaccumulation of heavy metals in organs of freshwater fish *Cyprinus carpio* (common carp). *International Journal of Environmental Science Technology*. 5(2): 179-182.
40. Wakeel, J., Ehsan, N., Akhtar, R. W., and Shah, S. A. H. (2020). Morphology, Histopathology, and hematology as biomarkers of cadmium toxicity in field rats. *Iranian Journal of Toxicology*. Jan;14(1):33-41.

41. Wang, H. J., Liu, Z. P., Jia, X. D., Chen, H., and Tan, Y. (2014). Endocrine disruption of cadmium in rats using the OECD enhanced TG 407 test system. *Biomedical Environment Sciences*. 27(12): 950-959.
42. Zhou, Z., Lu, Y., Pi, H., *et al.*(2016). Cadmium exposure is associated with the prevalence of dyslipidemia. *Cellular Physiology and Biochemistry*. 40: 633-643.

Access this Article in Online	
	Website: <a href="http://www.ijarbs.com">www.ijarbs.com</a>
	Subject: Biochemistry
Quick Response Code	
DOI: <a href="https://doi.org/10.22192/ijarbs.2021.08.08.001">10.22192/ijarbs.2021.08.08.001</a>	

How to cite this article:

Kshama Dwivedi. (2021). Hematological and Biochemical Investigations in Albino Rat Exposed to Cadmium Chloride. *Int. J. Adv. Res. Biol. Sci.* 8(8): 1-9.  
DOI: <http://dx.doi.org/10.22192/ijarbs.2021.08.08.001>