
International Journal of Advanced Research in Biological Sciences

ISSN : 2348-8069

www.ijarbs.com

Research Article



The Role of *Aloe vera* for the Protection Earthworms

Rajesh Kumar Singh*

Assistant professor, Department of Chemistry, Jagdam College, J P University, Chapra, India-841301.

*Corresponding author

Abstract

Earthworm is life of soils. It regulates soils productivity, humidity and concentration of water, oxygen, minerals, temperature and pH. The life of earthworm is getting destroyed by application of artificial insecticides, pesticides, herbicides, fertilizers, various types of wastes and acid rain. These killer substances change physical, chemical and biological properties of soils and generate corrosive atmosphere for earthworm. These pollutants change the concentration of natural minerals which are present in soils by increasing the acidic character of soils. The acidic soils develop microbiological corrosion cell with earthworm and it oxidize carbohydrate, protein and fat into CO₂, H₂O, NH₃, glycerol and organic acid. Corrosion reaction increases the temperatures of soils and finally kill earthworm. Heavy metals and gaseous pollutants come into soil by different medium which produce corrosive effect for earthworm. Pyrite ores of industrial area enter into soil to form acid that acid is corroding earthworm. These foreign materials contaminate soil and reduce the fertility of soil. Aloe Vera juice is applied to control the pH values of soils and protect the life of earthworm. The experimental work shows that Aloe Vera has capability to control the physical, chemical and biological parameters of soils and it also enhances the fertility and productivity of soil and protects the life of earthworm by corrosive pollutants. For this work soil samples are taken from the coal area of Jhariya district (in Dhanbad), steel plant area Chas in Bokaro district, urban area of Chak Beyriya in district of Patna and village area of Fulwariy-Tajpur in district of Chapra.

Keywords: Soils, Earthworm, biological corrosion cell, pollutants, pH, Aloe Vera.

Introduction

Earthworms are very important components of soils because they can regulate fertility of soils [1], its humidity [2] and pH [3]. But these organisms are destroyed [4] by interaction of industrial's pollutants, effluents, flues, hazardous wastes, municipal wastes, households wastes, hospital wastes, artificial fertilizers, pesticides, insecticides, herbicides, rodenticides, particulates, corrosive gases, acid rain and global warming. These substances are mostly acidic and basic in character and they can form a microbiological corrosion cell [5] with earthworms' thus biological corrosion reaction starts with these species and in this way their morphology can be changed which leads to the destruction of these organism in soils [6]. These harmful substances

change physical, chemical and biological properties of soils. They can disturb pH values and minerals composition of soils [7].

The major sources of corrosive pollutants, effluents, flues and hazardous wastes which contaminating soils are chemical industry, acid manufacture, sugar, coal mines, washery, coke manufacture, distillery, electroplating, paint manufacture, petroleum refinery, plastic manufacture, pulp and paper industry, steel industry, tannery, textile processing, electronic equipment, city waste, fertilizer industry, pesticides and herbicides industry, mining and ore processing, metallurgy, chemical industry, alloys, leather, electrical power plant, nuclear reactor, soap

and detergent industry, synthetic rubber, medicines, cosmetics, adhesives, explosives, salts, food processing, automobile industry, bricks making industry, rice and flour mills, glass and ceramic industry, cement industry.

The above mentioned industries release directly and indirectly harmful hazardous wastes into ponds, lakes, canals, rivers, sea and ocean and these pollutants can contaminate soils [8], atmosphere [9] and water sources [10]. Ponds, canals, rivers and ground water are used for irrigation of soils and this polluted water sources create corrosive environment for earthworms [11].

The inorganic pollutants are oxide of carbon, oxide of nitrogen, oxide of sulphur, oxide of halogen, hydride of sulphur, hydride of halogen, ammonia and organic pollutants are aldehyde (formaldehyde, acetaldehyde) and ketone (acetone), carboxylic acid (formic acid and acetic acid), pyridine, alcohol (methyl alcohol, ethyl alcohol), thiol (methyl thiol, ethyl thiol), methyl isocyanide, amines (methylamine and ethylamine) released by various industrial and nonindustrial sources. The concentrations of these pollutants are increased into atmosphere and they absorb moisture to convert into acids [12]. These acids condense into dewdrops and come into soils. These acids produce not only bad effects on soils and earthworms but also change the pH value of soils.

Acid rain can play important role for destruction of soils' organisms [13] and adversely affect pH of soils [14]. Soils possess natural minerals like Na, K, Fe, Cu, Zn, P, N₂, O₂ and the composition of these minerals are reduced by acid rain. The concentration of CO₂, CH₄ and water vapour are increased in atmosphere so the earth's temperature is raised and it creates problems for the survival of soils organisms.

Particulates come into atmosphere due to large scale industrialization, urbanization, deforestation, agriculturalization, infrastructure development works, mining and minerals processing works, stones breaking works, constructions works, making of railway tracks etc. These particulates contain heavy metals Fe, Cr, Co, Ni, Cu, Zn, Mo, As, Pb, Bi, and soot's of carbon, sulphur and fly ash. These metals are deposited on the surface of different components of environment and enter into soils directly and indirectly through various sources. These heavy metals

contaminate soil and alter soils physical, chemical and biological properties. They also develop hostile environment for soil organisms [15] and change its pH values.


Experimental

Collecting soils of different region like mining area (Jhariya in dist. of Dhanbad), industrial area (Chas in dist. of Bokaro), urban area (Chak Beyriya in dist of Patna) and village area (Fulwariy-Tajpur in dist of Chapra) and examine their pH values, concentration of minerals and oxygen and temperature. Earthworms are dispersed in different types of soils in the absence and presence of Aloe Vera and their physical, chemical and biological activities are observed at different concentrations of Aloe Vera, temperatures and times. The pH values of soils are recorded in different areas absence and presence of Aloe Vera at intervals of time. Earthworm's biological corrosion activities are calculated at 24hrs, 48hrs, 72hrs, 96hrs and 120hrs.


Results and Discussion

Analysis of pH values of soils in different regions and their values are mentioned in Table1. It is observed that mining area soils are more acidic with respect of other areas. The number of earthworms aided in various types of pH values of soils are counted at different intervals of time and the results are depicted in Table1 and figure2. Results of Table1 and figure2 indicated that mining area soil killed more earthworms. The acidic soils developed bio-electrochemical cell with earthworms and oxidized its amino acids and carbohydrates into ammonia, water and carbon dioxide. Fat dissociated in form glycerol and organic acids in presences of acids which oxidized into alkane and carbon dioxide. The bio-electrochemical cell occurs with earthworm and its mechanism as it is shown in figure1.

Anodic reaction with aminoacids




Cathodic reaction




Anodic reaction with carbohydrate



Cathodic reaction



Anodic reaction with Fats



Cathodic reaction





Figure1. Bioelectrochemical corrosion reaction with earthworm

Table1. pH values of soils without Aloe Vera in presence of earthworm

Soils of different regions (150g)	pH	Earthworm	24hrs	48hrs	72hr	96hrs	120hrs
Mining area (Jhariya in Dhanbad)	5.6	25	18	15	11	6	1
Industrial area (Chas,Bokaro)	6.2	25	20	17	14	12	9
Urban area (Chak Beriya)	6.4	25	22	19	16	13	10
Village area (Fulwariya, Chapra)	6.7	25	24	23	20	18	15





Figure2. Numbers of earthworm in different pH of soils Vs. time (hrs)

Bioelectrochemical reaction mechanism for earthworms indicated that they are corroded in acidic medium. Aloe Vera is used to control corrosive nature of soils. The solution of Aloe Vera was added into different pH values of soils at variation of times and its results were recorded in table 2 and figure 3. The results

of Table 2 indicated that pH values of soils increased after addition of Aloe Vera and it minimized biological corrosion reaction. It is observed that number of earthworms increased after addition of Aloe Vera. It suppressed the concentration of H⁺ ions and controlled formation of biological corrosion cell.

Table 2. pH values of soils with Aloe Vera in presence of earthworm

Soils of different regions (150g)	pH	No. of Earthworms	24hrs	48hrs	72hr	96hrs	120hrs
Mining area (Jhariya in Dhanbad)	6.0	25	24	27	29	32	35
Industrial area (Chas, Bokaro)	6.5	25	25	30	34	37	45
Urban area (Chak Beriya)	6.8	25	27	33	38	47	51
Village area (Fulwariya, Chapra)	6.9	25	28	37	42	50	65



Earthworm’s weight was measured in absence and presence of Aloe Vera into different pH values of soils and its results are depicted in Table 3 and figure4. It is noticed that its weight varies in different nature of soils. The weight of earthworm’s increased in Aloe

Vera mixed soils with respect of unaided soils. The figure 5 indicated that after addition of Aloe Vera earthworms improved their physical and biological properties.

Table 3. Earthworm position in different soils without and with Aloe Vera

Soils	Mining area	Industrial area	Urban area	Village area
pH	5.6	6.2	6.4	6.7
Wt. of earthworm(mg) without Aloe Vera	5.103	5.103	5.103	5.103
Wt. of earthworm (mg) without Aloe Vera after 24hrs	3.124	4.231	4.534	4.786
Wt. of earthworm (mg) with Aloe Vera after 24hrs	3.956	4.645	4.812	5.367






Figure 5. Earthworm kept in Aloe Vera aided soil

Heavy metals entered into soils by different sources and its compositions are recorded in Table4. They reacted with soils to acidic and basic compounds which generated corrosive atmosphere for earthworms and increased temperature of soils reducing humidity,

concentration of oxygen, water and others natural minerals. Heavy metals concentrations were analyzed after addition of Aloe Vera; it is observed that their concentrations were decreased after addition of Aloe Vera.

Table 4. Concentration of external minerals into Soils

Minerals (in PPM)	Mining area soils	Industrial area soils	Urban area soils	Village area soils
Fe	632	328	125	59
Ni	423	231	101	23
Cr	124	89	25	12
Pb	78	67	21	5
As	142	134	57	10
Zn	76	55	32	8
Cu	112	99	34	10
Mo	78	64	7	2
Bi	34	23	00	00
Aloe Vera (100ml) 120hrs	Mining area soils(250g)	Industrial area soils(250g)	Urban area soils (250g)	Village area soils (250g)
Fe	341	211	75	29
Ni	159	131	47	10
Cr	77	43	5	2
Pb	30	18	7	4
As	99	105	22	6
Zn	44	13	14	00
Cu	69	49	11	00
Mo	19	25	1	00
Bi	9	8	00	00

Conclusion

Earthworms are life of soils. Soil's physical, chemical and biological properties depend on availability of earthworms. Their availabilities are reducing by application of large scale artificial fertilizers, pesticides, insecticides, herbicides and rodents. The other pollutants like industrials effluents, household waste, biological wastes, municipals wastes, mining water, particulates and harmful chemical and gases come into soils directly and indirectly by different sources and contaminate soils destroying earthworms. These pollutants altered the pH values of soils and produced question mark on the survival of earthworm. Aloe Vera is used as remedy for soils by corrosive pollutants. The experimental results show that Aloe Vera has capability to control pH of soils, temperature, humidity, concentration of oxygen and composition of minerals and it creates ecofriendly atmosphere for soils and increasing the number of earthworms as well as their productivity.

Acknowledgments

Author is thankful to the UGC, New Delhi for providing financial support. Author is also thankful to the department of chemistry, Jagdam College, J P, University, Chapra for providing laboratory facilities.

References

1. Bloomfield JP, Williams RJ, Goody DC, Cape JC, Guha P (2006) Impact of climate change on the fate and behavior of pesticides in surface and groundwater, *Science of the Total Environment* 369, 163-177.
2. Baxter RA, Gilbert RE, Lidgett RA, Mainprize JH and Vodden HA (1975) The degradation of PCBs by microorganisms, *Sci. Toatal Environ.* 4, 53-61.
3. Balthazor TM and Hallas LE (1985) Glyphosate – degrading microorganisms from industrial activated sludge *Applied and Environmental Microbiology*, *Bol.* 51, 432-434.
4. Arias-Estevez M, Lopez-Periago E, Martinez-Carballo E (2008) (Review), The mobility and degradation of pesticides in soils and the pollution of groundwater resources. *Agriculture, Ecosystems and Environment* 123, 247-260. *Eurasian Soil Science* 39, 1271-1283.
5. Chen G, Abichou T, Subramaniam PK (2007), Impact on surface charge density on colloid deposition in unsaturated porous media, *Colloids and Surfaces A: Physicochem. Eng. Aspects* 302, 342-348.
6. Celis R, Hermosin MC, Carrizosa MJ, Cornejo J (2002) Inorganic and organic clays as carriers for controlled release of the herbicide hexazione, *J. Agric. Food Chem.* 50, 2324-2330.
7. Dekker LW, CJRitsema and K Oostindie (2004) Dry Spots in Golf Courses: Occurrence, Amelioration and Prevention, *Acta Horticulture* 661, 99-104.
8. Eggleton J and Thomas KV (2004) A review of factors affecting the release and bioavailability of contaminants during sediment disturbance events, *Environment International* 30, 973-980.
9. Eriksson E, Baun A, Mikkelesen PS and Ledin A (2007) Risk assessment of xenobiotics in stromwater discharged to Harrestrup A, Denmark *Desalination* 215, 187-197.
10. Feng JC, Thompson DG and Reynolds PE (1990) Fate of Glyphosate in a Canadian Forest Watershed, 1. Aquatic Residues and Off-Target Deposit Assessment, *J. Agric. Food Chem.* 38, 1110-1118.
11. Feng X, Simpson A, Simpson M (2006) Investigation the role of mineral-bound humic acid in phenanthrene sorption, *Environmental Science and Technology* 40, 3260-3266.
12. Ghanem A, Bados P, Estaum AR and Mougine C (2007) Concentrations and specific loads of glyphosate, diuron, atrazin, nonylphenol and metabolites thereof in French urban sewage sludge, *Chemosphere* 69, 1368-1373.
13. Gurses A, Karaca S, Yalcin M (2004) Determination of adsorptive properties of clay|water system :methylene blue sorption, *J. Colloid Interface Sci.* 269, 310-314.
14. Lombardi BM, Torres Sanchez RM, Eloy P, Genet M (2006) Interaction of thiabendazole and benzimidazole with montmorillonite, *Appl. Caly Sci.* 33, 59-65.
15. Wick LY, Shi L, Harms H (2007) Electro-bioremediation of hydrophobic organic soil contaminants: A review of fundamental interactions, *Electrochimica Acta* 52, 3441-3448.